arXiv:1209.3313v2 [cond-mat.dis-nn] 18 Sep 2012

Anomalous fluctuations of currents in Sinai-type random chains with strongly correlated disorder

Gleb Oshanin,[1, *] Alberto Rosso,[2, †] and Grégory Schehr[2, ‡]

[1] *Laboratoire de Physique Théorique de la Matière Condensée (UMR CNRS 7600), Université Pierre et Marie Curie (Paris 6) - 4 Place Jussieu, 75252 Paris, France*

[2] *Laboratoire Physique Théorique et Modèles Statistiques (UMR CNRS 8626), Université de Paris-Sud, Orsay Cedex, France*

(Dated: September 19, 2012)

We study properties of a random walk in a generalized Sinai model (SM), in which a quenched random potential is a trajectory of a *fractional* Brownian motion with arbitrary Hurst parameter H, $0 < H < 1$, so that the random force field displays strong spatial correlations. In this case, the disorder-average mean-square displacement (MSD) grows in proportion to $\log^{2/H}(n)$, n being time. We prove that moments of arbitrary order k of the steady-state current J_L through a finite segment of length L of such a chain decay as $L^{-(1-H)}$, independently of k, which suggests that despite a logarithmic confinement the average current is much higher than its Fickian counterpart in homogeneous systems. Our results reveal a paradoxical behavior such that, for fixed n and L, the MSD *decreases* when one varies H from 0 to 1, while the average current *increases*. This counter-intuitive behavior is explained via an analysis of representative realizations of disorder.

PACS numbers: 05.40.-a, 02.50.-r,05.10.Ln

Since the pioneering works [1–3], random walks in random environments have attracted a great deal of attention. In part, due to a general interest in dynamics in disordered systems, but also because such random walks have found many physical applications. Among them we mention dynamics of the boundary separating the coil and helix phases in a random heteropolymer [4, 5], dynamics in a random-field Ising model [6, 7], motion of dislocations in disordered crystals [8], mechanical unzipping of DNA [9], translocation of biomolecules through nanopores [10] and the dynamics of molecular motors [11]. Some functionals arising in this model, like probability currents in finite samples, show up in mathematical finance [12, 13]. Other examples can be found in [14–16].

In the discrete formulation, one considers a random walk on a discrete lattice evolving in a discrete time. At each time step the walker jumps from site X to site $X \pm 1$ with site-dependent probabilities p_X to $X+1$ and $q_X = 1 - p_X$ to $X-1$ given by $p_X = \frac{1}{2}(1 + \varepsilon \cdot s_X)$, where the amplitude $0 < \varepsilon < 1$ measures the strength of the disorder and s_X are independent and identically distributed (i.i.d.) random variables (r.v.). One often assumes binomial r.v., i.e., $s_X = \pm 1$ with probability p and $1 - p$. The probability $P_n(X)$ of finding the walker at site X after n time steps satisfies

$$P_n(X) = q_{X+1}\, P_{n-1}(X+1) + p_{X-1}\, P_{n-1}(X-1)\,. \quad (1)$$

In case of no global bias ($p = 1/2$), i.e. for the so-called Sinai model (SM), a remarkable result [3] is that for a given environment $\{p_X\}$ the squared displacement

$$X_n^2 \sim m(\{p_X\})\, \ln^4(n)\,, \quad (2)$$

as $n \to \infty$ with probability almost 1, where $m(\{p_X\})$ is a function of the environment only [17]. Another intriguing feature of the SM concerns transport properties. It was revealed by analysing the probability current J_L through a finite Sinai chain of length L, that the disorder-average current decays as $1/\sqrt{L}$ [18–21]. Curiously enough, despite a logarithmic confinement (2), the disorder-average current appears to be anomalously high, so that such disordered chains offer on average less resistance with respect to transport of particles than homogeneous chains (all $p_X \equiv 1/2$) for which one finds Fick's law $J_L \sim 1/L$.

It is well-known that random walk in uncorrelated random environment $\{p_X\}$ can be considered as a random walk in presence of a random potential V_L which represents itself a random walk in space. Indeed, on scales of order L a random walk "explores" the potential

$$V_L = \sum_{X=1}^{L-1} \ln\left(p_X/q_X\right) = \ln\left(\frac{1+\varepsilon}{1-\varepsilon}\right) \sum_{X=0}^{L-1} s_X\,, \quad (3)$$

which is, clearly, a trajectory of a random walk with step length $\sigma = \ln((1+\varepsilon)/(1-\varepsilon))$. The case of standard Sinai diffusion where the r.v. s_X are independent is now very well understood. On the contrary, there hardly exist analytical results in the case where these r.v. are strongly correlated. Such correlations are important, for instance, to understand the dynamics of the helix-coil boundary in random heteropolymers, where the location of chemical units usually exhibits strong correlations [22]. Such correlations are also currently studied in mathematical finance, as a necessary improvement over the standard Black-Sholes-Merton (BSM) model [23]. Any exact result for such situations would thus be welcome and here we make a step further in this direction.

In this Letter, we study properties of random walks in random environments in which the transition probabilities $\{p_X\}$ are strongly correlated so that the potential V_L in (3) is a fractional Brownian motion (fBm): V_L is

Gaussian, with $V_{L=0}=0$ and moments

$$\mathbb{E}\left\{V_L\right\}=0\,,\ \mathbb{E}\left\{(V_L-V_{L'})^2\right\}=\sigma^2|L-L'|^{2H}\,, \qquad (4)$$

where $\mathbb{E}\{\ldots\}$ here and henceforth denotes averaging over realizations of V_L and $0<H<1$. The case $H=1/2$ corresponds to the original SM. For $H<1/2$ the potential is subdiffusive while for $H>1/2$ it is superdiffusive.

The mean squared displacement $\mathbb{E}\{X_n^2\}$ in a correlated random environment can be estimated as follows. Assuming Arrhenius' law for the activated dynamics [14], the time t_L required for a particle to diffuse in a disordered potential V_L over a scale of order of L, is of order of $t_L\sim e^{V_L^*}$, where V_L^* is a typical energy barrier. For V_L in (4), $V_L^*\sim\sigma L^H$ for typical realizations of disorder, so that one finds that for sufficiently large times,

$$\mathbb{E}\left\{X_n^2\right\}\sim\sigma^{-2/H}\ln^{2/H}(n)\,. \qquad (5)$$

Our focal interest here is the understanding of the behavior of the disorder-average current J_L through a finite sample (of length L) of such a disordered chain, of its moments of arbitrary order, and eventually, of the full probability density function (pdf) of J_L. We proceed to show that, while its typical value is exponentially small $J_{\rm typ}\sim\exp\left(-L^H\right)$, its moments decay algebraically

$$\mu_k(L)\equiv\mathbb{E}\left\{\left(J_L\right)^k\right\}\sim A_kL^{-\theta}\ ,\ L\gg1\,, \qquad (6)$$

where $\theta=1-H$ is the persistence exponent of the fBm [24, 25] and A_k is a L-independent constant. We remind that the persistence exponent associated to a stochastic process characterizes the algebraic decay of its survival probability $S(t)\sim t^{-\theta}$ [26, 27].

The result in (6) is rather astonishing: it states first that the moments of the current of an arbitrary order decay in the same fashion. Second, one observes that for arbitrary H, $0<H<1$, the disorder-average current in such random chains is larger than the Fickian current in homogeneous systems. Finally, on comparing (5) and (6) for fixed n and L sufficiently large, and varying H, one concludes that $\mathbb{E}\{X_n^2\}$ *increases* when H goes from 1 to 0, while the disorder-average current *decreases*, which is an absolutely counter-intuitive and surprising behavior.

In what follows we prove (6) and explain this astonishing behavior using three different, complementary approaches: (i) a rigorous one, based on exact bounds, for the discrete random walk in a fBm potential, (ii) scaling arguments for the continuous version, both in space and time, of this diffusion (Langevin dynamics), which also allows to investigate the pdf of J_L (not only the moments) and (iii) thorough numerical simulations. We argue that (6) also holds for any disordered potential V_L in (3) which is the trajectory of a stochastic process with persistence exponent θ. As shown below, the occurrence of θ in (6) can be physically understood as the moments of J_L are dominated by the configurations of V_L which drift to $-\infty$ without re-crossing the origin (see also [28]). Such configurations occur with a probability $\sim L^{-\theta}$ (L playing the role of time), yielding the L-dependence in (6).



FIG. 1. (Color online) The disorder-average current $\mathbb{E}\left\{J_L\right\}$ (squares) and $\mu_k(L)$ with $k=2$ (circles) and $k=3$ (triangles) as a function of L for the fBm with $H=0.75$, $H=0.4$ and $H=0.25$ (from top to bottom). The solid line is the predicted behavior $\sim L^{-\theta}$ (6) with $\theta=1-H$. The temperature is $T=0.25$ and averaging is performed over 10^5 samples. We used arbitrary units [a.u.] because the data for each value of H have been vertically shifted for clarity.

We first consider the discrete version of SM, take a finite segment of length L and impose fixed concentrations of particles at the endpoints, $P_n(X=0)=P_0$ and $P_n(X=L)=P_L$. We then solve (1) for a fixed environment $\{p_X\}$, and find that the steady-state current of particles through the chain J_L is given by [18, 19]

$$J_L=\frac{D_0P_0}{\tau_L}-\frac{D_0P_L}{\tau_L^*}, \qquad (7)$$

where $D_0=1/2$ is the diffusion coefficient of a homogeneous chain, τ_L is the so-called Kesten variable [29]:

$$\tau_L=1+\frac{p_1}{q_1}+\frac{p_1p_2}{q_1q_2}+\ldots+\frac{p_1p_2\ldots p_{L-1}}{q_1q_2\ldots q_{L-1}}\,, \qquad (8)$$

and τ_L^* is obtained from (8) by replacements $p_k\to q_{L-k}$ and $q_k\to p_{L-k}$. Thinking of L as "time", one notices that τ_L and τ_L^* are time-averaged discretized geometric fractional Brownian motions (they can be thought of as the "prices" of Asian options within the framework of the fractional BSM model [13]). Note that in absence of a global bias $\mathbb{E}\{1/\tau_L\}=\mathbb{E}\{1/\tau_L^*\}$, and hence, without any loss of generality we set $P_L=0$ in what follows. Thus, combining (3, 7, 8) and setting $P_0=1$ yields

$$J_L=\frac{1}{2}\left(1+\sum_{l=1}^{L-1}\exp\left(V_l\right)\right)^{-1}\,. \qquad (9)$$

For typical realizations of $\{p_X\}$, the size of $|V_l|$ is $\mathcal{O}(l^H)$ so that the typical current $J_{\rm typ}$ is $J_{\rm typ}\sim\exp\left(-L^H\right)$. We now turn to the disorder-average current and establish upper and lower bounds on $\mu_k(L)$ (6).

To obtain an upper bound, consider a given realization of the sequence $V_1, V_2, \ldots, V_{L-1}$ and denote the maximal value among them as $V_{\max} = \max_{0 \le i \le L-1} V_i$. From (8) one has $\tau_L = (1 + \sum_{l=1}^{L-1} V_l) \ge \exp(V_{\max})$, so that

$$J_L^k \le (1/2)^k \exp(-k V_{\max}) \; . \qquad (10)$$

Hence, averaging in (10) amounts to calculating the characteristic function of the distribution of $V_{\max}$. Since $\exp(-kV_{\max})$ vanishes in the limit $L \to \infty$ (recall that $V_{\max} \sim L^H$) the average value of $\exp(-kV_{\max})$ is dominated by configurations of the potential with $V_{\max} \to 0$. The asymptotic behavior of the pdf $P_L(V_{\max})$ for fixed $V_{\max}$ and large L is known from [24, 25], yielding $\ln P_L(V_{\max}) = \theta \ln L^{-1} + \mathcal{O}(1)$, where $\theta = 1 - H$ is the persistence exponent [30]. Therefore, for $L \to \infty$ one has

$$\mu_k(L) \le B_k L^{H-1} \, , \qquad (11)$$

where B_k is an L-independent constant.

To determine a lower bound on $\mu_k(L)$ we follow [18, 19, 31] and make the following observation: averaging the expression in (9) amounts to taking average over the set Ω of all possible trajectories $\{V_l\}_{1 \le l \le L}$. Since $\tau_L > 0$, a lower bound on $\mu_k(L)$ can be straightforwardly obtained if one averages instead over some finite subset $\Omega' \subset \Omega$ of trajectories with some prescribed properties:

$$\mu_k(L) \ge \mathbb{E}_{\Omega'}\left\{J_L^k\right\} \; . \qquad (12)$$

We choose Ω' as comprising all possible trajectories $\{V_l\}_{0 \le l \le L}$ which, starting at the origin at $l = 0$, never cross the deterministic curve $Y_l = Y_0 - \alpha \ln(1 + l)$ with $Y_0 > 0$ and $\alpha > 1$. For any such trajectory

$$\tau_L = 1 + \sum_{l=1}^{L-1} \exp(V_l) \le \sum_{l=0}^{L-1} \exp(Y_l) \le \exp(Y_0)\zeta(\alpha) \, , \qquad (13)$$

where $\zeta(\alpha)$ is the zeta-function, which implies that

$$\mu_k(L) \ge \mathbb{E}_{\Omega'}\left\{J_L^k\right\} \ge (\exp(Y_0)\zeta(\alpha)/2)^{-k}\, \mathbb{E}_{\Omega'}\{1\} \; . \qquad (14)$$

One recognizes that $\mathbb{E}_{\Omega'}\{1\}$ is, by definition, the survival probability, S_L up to time L, for a fBm, starting at the origin in presence of a "moving trap" starting at Y_0 and evolving via $Y_l = Y_0 - \alpha \ln(1 + l)$. For standard Brownian motion ($H = 1/2$) in presence of a trap which moves as $-l^z$, the leading large-L behavior of the survival probability S_L is exactly the same as in the case of an immobile trap, provided that $z < 1/2$ [33]. It is thus physically plausible to suppose that the same behavior holds for a more general Gaussian process such as a fBm. That is, one expects that for any $H > 0$ the leading large-L behavior of $\mathbb{E}_{\Omega'}\{1\}$ will be exactly the same for an immobile trap and a logarithmically moving trap, i.e., that $S_L = \mathbb{E}_{\Omega'}\{1\} \sim Y_0^{\theta/H}/L^\theta$ as $L \to \infty$ [24, 25], where $\theta = 1 - H$ (and this can be shown rigorously, up to logarithmic corrections [28]). Consequently, we find

$$\mu_k(L) \ge D_k L^{H-1} \, , \qquad (15)$$

where D_k is independent of L. Note that the bounds in (10) and (15) show the same L-dependence and thus yields the exact result announced in (6).

We now turn to a continuous-time and -space description of the dynamics in a disordered fBm potential. The position of the particle at time t is defined by $x(t) \in [0, L]$ which obeys a Langevin equation

$$\dot{x} = -V'(x) + \eta(t) \, , \qquad (16)$$

where $V'(x)$ is a quenched random force [related to the hopping rates p_X in (1)] such that $V(x)$ is a fBm with Hurst exponent H (4) and $\eta(t)$ is a Gaussian thermal noise of zero mean and covariance $\langle \eta(t)\eta(t')\rangle = 2T\delta(t - t')$. The current J_L can be obtained from the Fokker-Planck equation corresponding to (16) and reads

$$J_L = T \left(\int_0^L \exp\left[V(x)/T\right] dx \right)^{-1} \, , \qquad (17)$$

[see (7, 9) with $D_0 = T$ (16) and $P_0 = 1$]. We remark here that the functional on the right-hand-side of (17) has been studied within a different context in [24, 32], and it has been shown that its first moment decreases as $L^{-\theta}$ with $\theta = 1 - H$. We focus next on the moments of higher order and on the pdf of the functional in (17).

Instead of J_L/T, which can be viewed as the inverse of the partition function of SM, we focus on the free energy $F = T \log(J_L/T)$ and study its pdf $\Pi_{T=0}(F)$. Let us first discuss the case $T = 0$ where $F = E_{\min} = \min_{0 \le x \le L} V(x)$. Recalling that $V(0) = 0$ (such that $E_{\min} < 0$), the cumulative distribution $q_L(E) = \Pr(E_{\min} > -E)$, with $E > 0$, coincides with the survival probability of $V(x)$ up to 'time' L starting in E at initial 'time' in presence of an absorbing boundary at $V = 0$. For fBm, and more generally for any self-affine process, $q_L(E)$ takes the scaling form $q_L(E) = Q(E/L^H)$. For $L \gg E^{1/H}$, $q_L(E)$ behaves algebraically with a persistence exponent θ [26], $q_L(E) \sim E^{\theta/H}/L^\theta$ ($\theta = 1 - H$ for fBm [24, 25]). In the opposite limit where $L \ll E^{1/H}$, $q_L(E)$ decays as a Gaussian [for Gaussian $V(x)$]. Therefore, one has for $\Pi_{T=0}(f) = \partial_E q_L(E)\big|_{E=-F}$

$$\Pi_{T=0}(F) = \begin{cases} 0 \, , \; F > 0 \\ L^{-\theta}|F|^{\theta/H-1} \, , \; -L^H \ll F < 0 \\ \exp(-F^2/2L^{2H}) \, , \; F \ll -L^H \; . \end{cases} \qquad (18)$$

What happens at finite T where $F = E - TS$ is now the balance between the energy E and the entropy S ? One expects the particle to be localized close to the minimum $E_{\min}$, which is of order $\mathcal{O}(L^H)$, while the maximum entropy $\sim \mathcal{O}(\ln L)$. Hence when $L \gg 1$ the main contribution to F comes from $E_{\min}$ so that for a given sample



FIG. 2. (Color online) Pdf of the free energy $-F/L^H$ for different system sizes: $L = 64, 256, 1024$ from bottom to top. The black curve represents the distribution of $E_{\min}/L^H$ for $L = 4096$. **Inset:** Pdf of $E_{\min} - F$ for different system sizes: $L = 64, 256, 1024$ from left to right. Histograms are computed using 10^5 samples and setting $T = 1$.

at finite T, F will be very close to $E_{\min}$. This is corroborated by our numerical simulations (see Fig. 2).

We now come back to the current distribution. Very small currents, $J_L \ll J_{\rm typ} \sim \exp(-L^H)$, correspond to $F \ll -L^H$ in (18) and one obtains that $P(J_L)$ is log-normal, $\ln P(J_L) \propto -\ln^2(J_L/T)$. Within the opposite limit, $J_L \gg J_{\rm typ}$, one finds from (18) that

$$P(J_L) \sim \frac{[\log(J_L/T)]^{\theta/H-1}}{J_L L^{\theta}} \,. \qquad (19)$$

This algebraic behavior holds up to a large cut-off value $J_{\max}$. At $T = 0$ we have a sharp cut-off at $J_{\max} = 1$ (18), at finite T, $P(J_L)$ has a fast decay which depends on the fluctuations of $V(x)$ at a short length scale close to the origin $x = 0$. For $L \gg 1$, the moments of the current are dominated by the regime where $J_{\rm typ} \ll J_L < J_{\max} \sim \mathcal{O}(1)$ (19) such that one gets $\mu_k(L) \sim 1/L^{\theta}$, (6).

To perform numerical simulations we consider a discrete random potential V_k, $k = 0, 1, \ldots L - 1$, which displays fBm correlations (4), we set $\sigma^2 = 1$. We use a powerful algorithm, as explained in [34, 35], which allows to generate very long samples of fBm paths. For each sample, we compute the current, $J_L = T[\sum_{k=0}^{L-1} \exp(-V_k/T)]^{-1}$, the free energy $F = T \ln(J_L/T)$ and the ground state energy $E_{\min} = \min_k V_k$. In Fig. 1 we show a plot of the first three moments as a function of L for different values of H. These plots show a very good agreement with our analytical predictions in (6). In Fig. 2, we show that the pdf of the rescaled free energy, F/L^H, at finite temperature T converges to the pdf of the rescaled ground state energy $E_{\min}/L^H$. The reason for this is that, for each sample, the difference between F and $E_{\min}$ grows very slowly with L, probably logarithmically (inset of Fig. 2). In the rescaled variables, this difference vanishes when $L \to \infty$.



FIG. 3. (Color online) The pdf in Eq. (21) for $\sigma^2 L^{2H} = 1/2$ (blue), $\sigma^2 L^{2H} = 2$ (red) and $\sigma^2 L^{2H} = 5$ (green).

We close this paper with an observation that such Sinai-type chains show an interesting transition to a diode-like behavior when σ (at fixed L) or L (at fixed σ) exceeds some critical values. To illustrate such a transition, consider a finite chain in which at site $X = 0$ we maintain a fixed concentration $P_0 = 1$ of one type of "white" particles and place a sink for them at $X = L$. On the other hand, at $X = L$ we introduce a source which maintains concentration 1 of "black" particles and place a sink for them at site $X = 0$. The particles are noninteracting and in the steady-state, for fixed $\{p_X\}$ we have two counter-currents of white (J_L^w) and black (J_L^b) particles. Obviously one has $\mathbb{E}\left\{(J_L^w)^k\right\} \equiv \mathbb{E}\left\{(J_L^b)^k\right\}$ for any $k > 0$. Consider next the random variable

$$\omega = J_L^w/(J_L^w + J_L^b) = \tau_L^*/(\tau_L^* + \tau_L) \,, \qquad (20)$$

which probes the likelihood of an event that for a given realization of disorder $J_L^w = J_L^b$. The pdf of ω can be calculated exactly to give

$$P(\omega) = \frac{1}{\sqrt{2\pi}\omega(1-\omega)\sigma L^H} \exp\left(-\frac{\ln^2\left(\frac{1-\omega}{\omega}\right)}{2\sigma^2 L^{2H}}\right) . \qquad (21)$$

A remarkable feature of (21) is depicted in Fig. 3: Namely, $P(\omega)$ changes modality when $\sigma^{1/H} L$ crosses a critical value $\sigma_c^{1/H} L_c = 2^{1/2H}$. For short chains (small disorder) $P(\omega)$ is unimodal and centered at $\omega = 1/2$: any given sample is transmitting particles in both directions equally well and, most probably, $J_L^w = J_L^b$. On contrary, for sufficiently strong disorder and/or long chain, $P(\omega)$ becomes bimodal with a local minimum at $\omega = 1/2$ and two maxima close to 0 and 1. This means that a given sample is most likely permeable only in one direction.

GO is partially supported by the ESF Research Network "Exploring the Physics of Small Devices", AR - by ANR grant 09-BLAN-0097-02 and GS - by ANR grant 2011-BS04- 013-01 WALKMAT. This project was partially supported by the Indo-French Centre for the Promotion of Advanced Research under Project 4604-3.

---

[*] oshanin@lptmc.jussieu.fr
[†] alberto.rosso@lptms.u-psud.fr
[‡] gregory.schehr@lptms.u-psud.fr

[1] H. Kesten, M. V. Kozlov and F. Spitzer, Compos. Math. **30**, 145 (1975).
[2] B. Derrida and Y. Pomeau, Phys. Rev. Lett. **48**, 627 (1982).
[3] Ya. G. Sinai, Theor. Probab. Appl. **27** 256, (1982).
[4] P. G. de Gennes, J. Stat. Phys. **12**, 463 (1975).
[5] G. Oshanin and S. Redner, EPL **85**, 10008 (2009).
[6] R. Bruinsma and G. Aeppli, Phys. Rev. Lett. **52**, 1547 (1984).
[7] T. Nattermann and J. Vilain, Phase Transit. **11**, 5 (1988).
[8] J. P. Harth and J. Lothe, *The theory of dislocations* (New York, McGraw-Hill, 1968).
[9] J.-C. Walter, A. Ferrantini, E. Carlon and C. Vanderzande, Phys. Rev. E **85**, 031120 (2012).
[10] J. Mathe et al., Biophys. J. **87**, 3205 (2004).
[11] Y. Kafri, D. K. Lubensky and D. R. Nelson, Biophys. J. **86**, 3373 (2004).
[12] A. Comtet, C. Monthus and M. Yor, J. Appl. Prob. **35**, 255 (1998).
[13] G. Oshanin and G. Schehr, Quant. Finance **12**, 1325 (2012).
[14] J. P. Bouchaud, A. Comtet, A. Georges, and P. Le Doussal, Ann. Phys. **201**, 285 (1990).
[15] B. D. Hughes, *Random walks and random environments* Vol.**2** (Clarendon Press, Oxford, 1995).
[16] M. Sheinman, O. Bénichou, R. Voituriez and Y. Kafri, J. Stat. Mech. P09005 (2010).
[17] the probability being taken over all random walks of n steps, starting at the same point.
[18] S. F. Burlatsky, G. Oshanin, A. Mogutov and M. Moreau, Phys. Rev. A **45**, R6955 (1992).
[19] G. Oshanin, S. F. Burlatsky, M. Moreau and B. Gaveau, Chem. Phys. **177**, 803 (1993).
[20] G. Oshanin, A. Mogutov and M. Moreau, J. Stat. Phys. **73**, 379 (1993).
[21] C. Monthus and A. Comtet, J. Phys. I France **4**, 635 (1994).
[22] E. N. Govorun et al., Phys. Rev. E **64**, 040903 (2001).
[23] C. Bender, T. Sottinen and E. Valkeila, in *Advanced mathematical methods for finance*, (ed. G. Di Nunno and B. Oksendal. Springer, Berlin), pp. 75 (2011), arXiv:1004.3106
[24] G. M. Molchan, Comm. Math. Phys. **205**, 97 (1999).
[25] J. Krug et al., Phys. Rev. E **56**, 2702 (1997).
[26] S. N. Majumdar, Current Science **77**, 370 (1999).
[27] F. Aurzada and T. Simon, preprint arXiv:1203.6554
[28] F. Aurzada, Elect. Comm. in Probab. **16**, 392 (2011).
[29] H. Kesten, Acta Math. **131**, 207 (1973).
[30] As a matter of fact, this result can be made more precise by writing the distribution $P_L(V_{\max})$ in a scaled form $P_L(V_{\max}) = L^{-H} f(y = V_{\max}/L^H)$. This implies that actually $P_L(V_{\max}) \sim V_{\max}^{(1-2H)/H}/L^{1-H}$ as $L \to \infty$.
[31] C. Monthus, G. Oshanin, A. Comtet, and S. F. Burlatsky, Phys. Rev. E **54**, 231 (1996).
[32] S. N. Majumdar, A. Rosso and A. Zoia, Phys. Rev. Lett. **102**, 120602 (2009).
[33] K. Uchiyama, Z. Wahrsch. Verw. Gebiete **54**, 75 (1980); P. L. Krapivsky and S. Redner, Am. J. Phys. **64**, 546 (1996).
[34] K. J. Wiese, S. N. Majumdar, A. Rosso, Phys. Rev. E **83**, 061141 (2011).
[35] R. Garcia-Garcia, A. Rosso and G. Schehr, Phys. Rev. E **81**, 010102(R) (2010)